

Time-dependent density-functional formalism

(for time being: electrons only, nuclei are fixed or treated classically)

(E. Runge, E.K.U.G., PRL 52, 997 (1984))

Basic 1-1 correspondence:

$v(\mathbf{r}t) \xleftrightarrow{1-1} \rho(\mathbf{r}t)$ The time-dependent density determines uniquely the time-dependent external potential and hence all physical observables for fixed initial state.

KS theorem:

The time-dependent density of the interacting system of interest can be calculated as density

$$\rho(\mathbf{r}t) = \sum_{j=1}^N \left| \varphi_j(\mathbf{r}t) \right|^2$$

of an auxiliary non-interacting (KS) system

$$i\hbar \frac{\partial}{\partial t} \varphi_j(\mathbf{r}t) = \left(-\frac{\hbar^2 \nabla^2}{2m} + v_s[\rho](\mathbf{r}t) \right) \varphi_j(\mathbf{r}t)$$

with the local potential


$$v_s[\rho(\mathbf{r}'t')](\mathbf{r}t) = v(\mathbf{r}t) + \int d^3r' \frac{\rho(\mathbf{r}'t')}{|\mathbf{r}-\mathbf{r}'|} + v_{xc}[\rho(\mathbf{r}'t')](\mathbf{r}t)$$

Proof of basic 1-1 correspondence between $v(\vec{r}t)$ and $\rho(\vec{r}t)$

define maps

$$F: v(\vec{r}t) \mapsto \Psi(t)$$

$$\tilde{F}: \Psi(t) \mapsto \rho(\vec{r}t)$$



$$G: v(\vec{r}t) \mapsto \rho(\vec{r}t)$$

The TDKS equations follow (like in the static case) from:

- i. the basic 1-1 mapping and**
- ii. the TD V -representability theorem (R. van Leeuwen, PRL 82, 3863 (1999)).**

A TDDFT variational principle exists as well, but this is more tricky (R. van Leeuwen, PRL 80, 1280 (1998)).

complete 1 - 1 correspondence not to be expected!

$$i \frac{\partial}{\partial t} \Psi(t) = \left(\hat{T} + \underline{\hat{V}(t)} + \hat{W} \right) \Psi(t) \quad \Psi(t_0) = \Psi_0$$

$$i \frac{\partial}{\partial t} \Psi'(t) = \left(\hat{T} + \underline{\hat{V}'(t)} + \hat{W} \right) \Psi'(t) \quad \Psi'(t_0) = \Psi_0$$

$$\hat{V}'(t) = \hat{V}(t) + C(t) \Leftrightarrow \Psi'(t) = e^{-i\alpha(t)} \Psi(t)$$

↑
“no operator”

with

$$\dot{\alpha}(t) = C(t)$$

$$\Rightarrow \underline{\underline{\rho'(\vec{r}t) = \rho(\vec{r}t)}}$$

$$\text{i.e. } \{ \hat{V}(t) + C(t) \} \rightarrow \rho(\vec{r}t)$$

If G invertible up to within time-dependent function $C(t)$

$\Rightarrow \Psi = FG^{-1}\rho$ fixed up to within time-dependent phase

i.e. $\Psi = e^{-i\alpha(t)}\Psi[\rho]$

For any observable \hat{O}

$$\langle \Psi | \hat{O} | \Psi \rangle = \langle \Psi[\rho] | \hat{O} | \Psi[\rho] \rangle = O[\rho]$$

is functional of the density

THEOREM (time-dependent analogue of Hohenberg-Kohn theorem)


The map

$$G : v(\vec{r}t) \mapsto \rho(\vec{r}t)$$

defined for all single-particle potentials $v(\vec{r}t)$ which can be expanded into a Taylor series with respect to the time coordinate around t_0

is invertible up to within an additive merely time-dependent function in the potential.

Proof:
to be shown:



i.e. $\hat{v}(\vec{r}t) \neq \hat{v}'(\vec{r}t) + c(t) \stackrel{!}{\Rightarrow} \rho(\vec{r}t) \neq \rho'(\vec{r}t)$

potential expandable into Taylor series

$$\exists k \geq 0 : \frac{\partial^k}{\partial t^k} [v(\vec{r}t) - v'(\vec{r}t)]_{t=t_0} \neq \text{constant}$$

step 1

$$\vec{j}(\vec{r}t) \neq \vec{j}'(\vec{r}t)$$

step 2

$$\rho(\vec{r}t) \neq \rho'(\vec{r}t)$$

Step 1: Current densities

$$\vec{j}(\vec{r}t) = \left\langle \Psi(t) \left| \hat{j}(\vec{r}) \right| \Psi(t) \right\rangle$$

$$\text{with } \hat{j}(\vec{r}) = -\frac{1}{2i} \sum_s \left(\left[\vec{\nabla} \hat{\psi}_s^+(\vec{r}) \right] \hat{\psi}_s(\vec{r}) - \hat{\psi}_s^+(\vec{r}) \left[\vec{\nabla} \hat{\psi}_s(\vec{r}) \right] \right)$$

Use equation of motion:

$$i \frac{\partial}{\partial t} \left\langle \Psi(t) \left| \hat{O}(t) \right| \Psi(t) \right\rangle = \left\langle \Psi(t) \left| i \frac{\partial}{\partial t} \hat{O}(t) + \left[\hat{O}(t), \hat{H}(t) \right] \right| \Psi(t) \right\rangle$$

$$\Rightarrow i \frac{\partial}{\partial t} \vec{j}(\vec{r}t) = \left\langle \Psi(t) \left[\hat{j}(\vec{r}), \hat{H}(t) \right] \right| \Psi(t) \right\rangle$$

$$i \frac{\partial}{\partial t} \vec{j}'(\vec{r}t) = \left\langle \Psi'(t) \left[\hat{j}(\vec{r}), \hat{H}'(t) \right] \right| \Psi'(t) \right\rangle$$


note: $\vec{j}(\vec{r}\underline{t}_0) = \vec{j}'(\vec{r}\underline{t}_0) = \left\langle \Psi_0 \left| \hat{j}(\vec{r}) \right| \Psi_0 \right\rangle \equiv \vec{j}_0(\vec{r})$

$$\rho(\vec{r}\underline{t}_0) = \rho'(\vec{r}\underline{t}_0) = \left\langle \Psi_0 \left| \hat{\rho}(\vec{r}) \right| \Psi_0 \right\rangle \equiv \rho_0(\vec{r})$$

$$\begin{aligned}
i \frac{\partial}{\partial t} [\vec{j}(\vec{r}t) - \vec{j}'(\vec{r}t)]_{t=t_0} &= \left\langle \Psi_0 \left[\left[\hat{j}(\vec{r}), \hat{H}(t_0) - \hat{H}'(t_0) \right] \right] \Psi_0 \right\rangle \\
&= \left\langle \Psi_0 \left[\left[\hat{j}(\vec{r}), V(t_0) - V'(t_0) \right] \right] \Psi_0 \right\rangle \\
&= i \rho_0(\vec{r}) \vec{\nabla} (v(\vec{r}t_0) - v'(\vec{r}t_0))
\end{aligned}$$

if $\frac{\partial^k}{\partial t^k} [v(\vec{r}t) - v'(\vec{r}t)]_{t=t_0} \neq \text{constant}$ holds for $k=0$

then $i \frac{\partial}{\partial t} [\vec{j}(\vec{r}t) - \vec{j}'(\vec{r}t)]_{t=t_0} \neq 0$



$\Rightarrow \underline{\underline{\vec{j}(\vec{r}t) \neq \vec{j}'(\vec{r}t)}} \quad \text{q.e.d.}$

if $\frac{\partial^k}{\partial t^k} [\vec{v}(\vec{r}t) - \vec{v}'(\vec{r}t)]_{t=t_0} \neq \text{constant}$ holds for $k > 0$

→ use equation of motion $k+1$ times:

$$\begin{aligned} \left(i \frac{\partial}{\partial t} \right)^2 \vec{j}(\vec{r}t) &= i \frac{\partial}{\partial t} \left\langle \Psi(t) \left[\hat{\vec{j}}, \hat{H}(t) \right] \Psi(t) \right\rangle \\ &= \left\langle \Psi(t) \left| i \frac{\partial}{\partial t} \left[\hat{\vec{j}}, \hat{H}(t) \right] + \left[\left[\hat{\vec{j}}, \hat{H}(t) \right], \hat{H}(t) \right] \right| \Psi(t) \right\rangle \end{aligned}$$

$$\begin{aligned} \left(i \frac{\partial}{\partial t} \right)^3 \vec{j}(\vec{r}t) &= i \frac{\partial}{\partial t} \left\langle \Psi(t) \left| i \frac{\partial}{\partial t} \left[\hat{\vec{j}}, \hat{H}(t) \right] + \left[\left[\hat{\vec{j}}, \hat{H}(t) \right], \hat{H}(t) \right] \right| \Psi(t) \right\rangle \\ &= \dots \end{aligned}$$

$$\left(i \frac{\partial}{\partial t} \right)^{k+1} \left[\vec{j}(\vec{r}t) - \vec{j}'(\vec{r}t) \right]_{t=t_0} = i \rho_0(\vec{r}) \vec{\nabla} \underbrace{\left(\left(i \frac{\partial}{\partial t} \right)^k \left[\vec{v}(\vec{r}t) - \vec{v}'(\vec{r}t) \right]_{t_0} \right)}_{\neq \text{constant}} \neq 0$$

⇒ $\vec{j}(\vec{r}t) \neq \vec{j}'(\vec{r}t)$ q.e.d.

Step 2: densities

Use continuity equation:

$$\frac{\partial}{\partial t} [\rho(\vec{r}t) - \rho'(\vec{r}t)] = -\text{div} [\vec{j}(\vec{r}t) - \vec{j}'(\vec{r}t)]$$

$$\begin{aligned} \Rightarrow \frac{\partial^{k+2}}{\partial t^{k+2}} [\rho(\vec{r}t) - \rho'(\vec{r}t)]_{t=t_0} &= -\text{div} \frac{\partial^{k+1}}{\partial t^{k+1}} [\vec{j}(\vec{r}t) - \vec{j}'(\vec{r}t)]_{t=t_0} \\ &= -\text{div} \rho_o(\vec{r}) \underbrace{\vec{\nabla} \left(\frac{\partial^k}{\partial t^k} [\mathbf{v}(\vec{r}t) - \mathbf{v}'(\vec{r}t)]_{t=t_0} \right)}_{\neq \text{constant}} \end{aligned}$$

remains to be shown:

$$\text{div} [\rho_o(\vec{r}) \vec{\nabla} u(\vec{r})] \neq 0 \quad \text{if} \quad u(\vec{r}) \neq \text{constant}$$

Proof: by reductio ad absurdum

Assume: $\operatorname{div}[\rho_o(\vec{r})\vec{\nabla}u(\vec{r})] = 0$ with $u(\vec{r}) \neq \text{constant}$

$$\begin{aligned} & \int d\vec{r}^3 \rho_o(\vec{r}) (\vec{\nabla}u(\vec{r}))^2 \\ &= -\int d\vec{r}^3 u(\vec{r}) \underbrace{\operatorname{div}[\rho_o(\vec{r})\vec{\nabla}u(\vec{r})]}_0 + \underbrace{\oint \rho_o(\vec{r})u(\vec{r})\vec{\nabla}u(\vec{r}) \cdot d\vec{S}}_0 = 0 \end{aligned}$$

$$\Rightarrow \rho_o(\vec{r})(\vec{\nabla}u(\vec{r}))^2 \equiv 0 \quad \longrightarrow \quad \begin{array}{l} \text{contradiction to} \\ u(\vec{r}) \neq \text{constant} \end{array}$$

Simplest possible approximation for $v_{xc}[\rho](\vec{r}t)$

Adiabatic Local Density Approximation (ALDA)

$$V_{xc}^{\text{ALDA}}(\vec{r}t) := V_{xc,\text{stat}}^{\text{hom}}(\mathbf{n}) \Big|_{\mathbf{n}=\rho(\vec{r}t)}$$

$V_{xc,\text{stat}}^{\text{hom}}$ = **xc potential of static homogeneous e-gas**

**Approximation with correct asymptotic $-1/r$ behavior:
time-dependent optimized effective potential**

(C. A. Ullrich, U. Gossmann, E.K.U.G., PRL 74, 872 (1995))

LINEAR RESPONSE THEORY

$t = t_0$: Interacting system in ground state of potential $v_0(\mathbf{r})$ with density $\rho_0(\mathbf{r})$

$t > t_0$: Switch on perturbation $\mathbf{v}_1(\mathbf{r}, t)$ (with $\mathbf{v}_1(\mathbf{r}, t_0) = 0$).

Density: $\rho(\mathbf{r}, t) = \rho_0(\mathbf{r}) + \delta\rho(\mathbf{r}, t)$

Consider functional $\rho[v](\mathbf{r}, t)$ defined by solution of interacting TDSE

Functional Taylor expansion of $\rho[v]$ around v_0 :

$$\begin{aligned}
 \rho[v](\mathbf{r}, t) &= \rho[v_0 + \mathbf{v}_1](\mathbf{r}, t) \\
 &= \rho[v_0](\mathbf{r}, t) && \longrightarrow \rho_0(\mathbf{r}) \\
 &\quad + \int \frac{\delta\rho[v](\mathbf{r}, t)}{\delta v(\mathbf{r}', t')} \Big|_{v_0} \mathbf{v}_1(\mathbf{r}', t') d^3r' dt' && \longrightarrow \rho_1(\mathbf{r}, t) \\
 &\quad + \frac{1}{2} \iint \frac{\delta^2\rho[v](\mathbf{r}, t)}{\delta v(\mathbf{r}', t') \delta v(\mathbf{r}'', t'')} \Big|_{v_0} \mathbf{v}_1(\mathbf{r}', t') \mathbf{v}_1(\mathbf{r}'', t'') d^3r' d^3r'' dt' dt'' && \longrightarrow \rho_2(\mathbf{r}, t) \\
 &\quad \vdots
 \end{aligned}$$

$\rho_1(\mathbf{r},t)$ = linear density response of interacting system

$$\chi(\mathbf{r}t, \mathbf{r}'t') := \left. \frac{\delta\rho[\mathbf{v}](\mathbf{r}t)}{\delta v(\mathbf{r}'t')} \right|_{\mathbf{v}_0} = \text{density-density response function of interacting system}$$


Analogous function $\rho_s[\mathbf{v}_s](\mathbf{r}t)$ for non-interacting system

$$\rho_s[\mathbf{v}_s](\mathbf{r}t) = \rho_s[\mathbf{v}_{s,0} + \mathbf{v}_{s,1}](\mathbf{r}t) = \rho_s[\mathbf{v}_{s,0}](\mathbf{r}t) + \int \left. \frac{\delta\rho_s[\mathbf{v}_s](\mathbf{r}t)}{\delta v_s(\mathbf{r}'t')} \right|_{\mathbf{v}_{s,0}} \mathbf{v}_{s,1}(\mathbf{r}'t') d^3r'dt' + \dots$$

$$\chi_s(\mathbf{r}t, \mathbf{r}'t') := \left. \frac{\delta\rho_s[\mathbf{v}_s](\mathbf{r}t)}{\delta v_s(\mathbf{r}'t')} \right|_{\mathbf{v}_{s,0}} = \text{density-density response function of non-interacting system}$$

GOAL: Find a way to calculate $\rho_1(\mathbf{r}, t)$ without explicitly evaluating $\chi(\mathbf{r}, t, \mathbf{r}', t')$ of the interacting system

starting point: Definition of xc potential

$$v_{xc}[\rho](\mathbf{r}, t) := v_S[\rho](\mathbf{r}, t) - v_{\text{ext}}[\rho](\mathbf{r}, t) - v_H[\rho](\mathbf{r}, t)$$


Notes:

- v_{xc} is well-defined through non-interacting/ interacting 1-1 mapping.
- $v_S[\rho]$ depends on initial determinant Φ_0 .
- $v_{\text{ext}}[\rho]$ depends on initial many-body state Ψ_0 .

\Rightarrow In general, $v_{xc} = v_{xc}[\rho, \Phi_0, \Psi_0]$

only if system is initially in ground-state then, via HK, Φ_0 and Ψ_0 are determined by ρ_0 and v_{xc} depends on ρ alone.

$$\left. \frac{\delta v_{\text{xc}}[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} = \left. \frac{\delta v_{\text{S}}[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} - \left. \frac{\delta v_{\text{ext}}[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} - \frac{\delta(t-t')}{|\mathbf{r}-\mathbf{r}'|}$$

$$\begin{array}{cccc}
\left. \frac{\delta v_{\text{xc}}[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} & = & \left. \frac{\delta v_{\text{S}}[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} & - & \left. \frac{\delta v_{\text{ext}}[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} & - & \frac{\delta(t-t')}{|\mathbf{r}-\mathbf{r}'|} \\
\uparrow & & \uparrow & & \uparrow & & \uparrow \\
\mathbf{f}_{\text{xc}}(\mathbf{r}t, \mathbf{r}'t') & & \chi_{\text{S}}^{-1}(\mathbf{r}t, \mathbf{r}'t') & & \chi^{-1}(\mathbf{r}t, \mathbf{r}'t') & & \mathbf{W}_{\text{C}}(\mathbf{r}t, \mathbf{r}'t')
\end{array}$$

$$\begin{array}{cccc}
\left. \frac{\delta v_{\text{xc}}[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} & = & \left. \frac{\delta v_{\text{S}}[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} & - & \left. \frac{\delta v_{\text{ext}}[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} & - & \frac{\delta(t-t')}{|\mathbf{r}-\mathbf{r}'|} \\
\uparrow & & \uparrow & & \uparrow & & \uparrow \\
\mathbf{f}_{\text{xc}}(\mathbf{r}t, \mathbf{r}'t') & & \chi_{\text{S}}^{-1}(\mathbf{r}t, \mathbf{r}'t') & & \chi^{-1}(\mathbf{r}t, \mathbf{r}'t') & & \mathbf{W}_{\text{C}}(\mathbf{r}t, \mathbf{r}'t')
\end{array}$$

$$\mathbf{f}_{\text{xc}} + \mathbf{W}_{\text{C}} = \chi_{\text{S}}^{-1} - \chi^{-1}$$

$$\begin{array}{cccc}
\left. \frac{\delta v_{\text{xc}}[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} & = & \left. \frac{\delta v_{\text{S}}[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} & - & \left. \frac{\delta v_{\text{ext}}[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} & - & \frac{\delta(t-t')}{|\mathbf{r}-\mathbf{r}'|} \\
\uparrow & & \uparrow & & \uparrow & & \uparrow \\
\mathbf{f}_{\text{xc}}(\mathbf{r}t, \mathbf{r}'t') & & \chi_{\text{S}}^{-1}(\mathbf{r}t, \mathbf{r}'t') & & \chi^{-1}(\mathbf{r}t, \mathbf{r}'t') & & \mathbf{W}_{\text{C}}(\mathbf{r}t, \mathbf{r}'t')
\end{array}$$

$$\chi_{\text{S}} \cdot \left| \mathbf{f}_{\text{xc}} + \mathbf{W}_{\text{C}} = \chi_{\text{S}}^{-1} - \chi^{-1} \right| \cdot \chi$$

$$\begin{array}{cccc}
\left. \frac{\delta v_{xc}[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} & = & \left. \frac{\delta v_S[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} & - & \left. \frac{\delta v_{ext}[\rho](\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} & - & \frac{\delta(t-t')}{|\mathbf{r}-\mathbf{r}'|} \\
\uparrow & & \uparrow & & \uparrow & & \uparrow \\
\mathbf{f}_{xc}(\mathbf{r}t, \mathbf{r}'t') & & \chi_S^{-1}(\mathbf{r}t, \mathbf{r}'t') & & \chi^{-1}(\mathbf{r}t, \mathbf{r}'t') & & \mathbf{W}_C(\mathbf{r}t, \mathbf{r}'t')
\end{array}$$

$$\chi_S \cdot \left| \mathbf{f}_{xc} + \mathbf{W}_C = \chi_S^{-1} - \chi^{-1} \right| \cdot \chi$$

$$\chi_S (\mathbf{f}_{xc} + \mathbf{W}_C) \chi = \chi - \chi_S$$

$$\chi = \chi_S + \chi_S (W_C + f_{xc}) \chi$$

Act with this operator equation on arbitrary $v_1(\mathbf{r}, t)$ and use $\chi v_1 = \rho_1$:

$$\rho_1(\mathbf{r}, t) = \int d^3 r' dt' \chi_S(\mathbf{r}, t, \mathbf{r}', t') \left[v_1(\mathbf{r}', t') + \int d^3 r'' dt'' \{ W_C(\mathbf{r}', t', \mathbf{r}'', t'') + f_{xc}(\mathbf{r}', t', \mathbf{r}'', t'') \} \rho_1(\mathbf{r}'', t'') \right]$$

- Exact integral equation for $\rho_1(\mathbf{r}, t)$, to be solved iteratively

- Need approximation for $f_{xc}(\mathbf{r}', t', \mathbf{r}'', t'') = \frac{\delta v_{xc}[\rho](\mathbf{r}', t')}{\delta \rho(\mathbf{r}'', t'')} \Big|_{\rho_0}$
(either for f_{xc} directly or for v_{xc})

Adiabatic approximation


$$v_{xc}^{\text{adiab}}[\rho](\mathbf{r}t) := v_{xc}^{\text{static DFT}}[\rho(\mathbf{t})](\mathbf{r}t)$$

In the adiabatic approximation, the xc potential $v_{xc}(\mathbf{r}t)$ at time t only depends on the density $\rho(\mathbf{r}t)$ at the very same point in time.

e.g. adiabatic LDA: $v_{xc}^{\text{ALDA}}(\mathbf{r}t) := v_{xc}^{\text{LDA}}(\rho(\mathbf{r}t)) = -\alpha \rho(\mathbf{r}t)^{1/3} + \dots$



$$\begin{aligned} \Rightarrow f_{xc}^{\text{ALDA}}(\mathbf{r}t, \mathbf{r}'t') &= \left. \frac{\delta v_{xc}^{\text{ALDA}}(\mathbf{r}t)}{\delta \rho(\mathbf{r}'t')} \right|_{\rho_0} = \delta(\mathbf{r} - \mathbf{r}') \delta(t - t') \left. \frac{\partial v_{xc}^{\text{ALDA}}}{\partial \rho(\mathbf{r})} \right|_{\rho_0(\mathbf{r})} \\ &= \delta(\mathbf{r} - \mathbf{r}') \delta(t - t') \left. \frac{\partial^2 e_{xc}^{\text{hom}}}{\partial n^2} \right|_{\rho_0(\mathbf{r})} \end{aligned}$$

Total photoabsorption cross section of the Xe atom versus photon energy in the vicinity of the 4d threshold.



Solid line: self-consistent time-dependent KS calculation [A. Zangwill and P. Soven, PRA 21, 1561 (1980)]; **crosses:** experimental data [R. Haensel, G. Keitel, P. Schreiber, and C. Kunz, Phys. Rev. 188, 1375 (1969)].

Photo-absorption in weak lasers



No absorption if $\omega < \text{lowest excitation energy}$

Standard linear response formalism

$H(t_0)$ = full static Hamiltonian at t_0

$$H(t_0)|m\rangle = E_m|m\rangle \quad \leftarrow \text{exact many-body eigenfunctions and energies of system}$$

full response function

$$\chi(r, r'; \omega) = \lim_{\eta \rightarrow 0^+} \sum_m \left(\frac{\langle 0 | \hat{\rho}(r) | m \rangle \langle m | \hat{\rho}(r) | 0 \rangle}{\omega - (E_m - E_0) + i\eta} - \frac{\langle 0 | \hat{\rho}(r') | m \rangle \langle m | \hat{\rho}(r') | 0 \rangle}{\omega + (E_m - E_0) + i\eta} \right)$$

\Rightarrow The exact linear density response

$$\rho_1(\omega) = \hat{\chi}(\omega) v_1(\omega)$$

has poles at the exact excitation energies $\Omega = E_m - E_0$

Discrete excitation energies from TDDFT

exact representation of linear density response:

$$\rho_1(\omega) = \hat{\chi}_S(\omega) \left(v_1(\omega) + \hat{W}_C \rho_1(\omega) + \hat{f}_{xc}(\omega) \rho_1(\omega) \right)$$

“ $\hat{}$ ” denotes integral operators, i.e. $\hat{f}_{xc} \rho_1 \equiv \int f_{xc}(\vec{r}, \vec{r}') \rho_1(\vec{r}') d^3 r'$

$$\text{where } \hat{\chi}_S(\vec{r}, \vec{r}'; \omega) = \sum_{j,k} \frac{M_{jk}(\vec{r}, \vec{r}')}{\omega - (\epsilon_j - \epsilon_k) + i\eta}$$

$$\text{with } M_{jk}(\vec{r}, \vec{r}') = (f_k - f_j) \varphi_j(\vec{r}) \varphi_j^*(\vec{r}') \varphi_k(\vec{r}') \varphi_k^*(\vec{r})$$

$$f_m = \begin{cases} 1 & \text{if } \varphi_m \text{ is occupied in KS ground state} \\ 0 & \text{if } \varphi_m \text{ is unoccupied in KS ground state} \end{cases}$$

$\epsilon_j - \epsilon_k$ KS excitation energy

$$\left(\hat{1} - \hat{\chi}_S(\omega) \left[\hat{W}_C + \hat{f}_{xc}(\omega) \right] \right) \rho_1(\omega) = \hat{\chi}_S(\omega) v_1(\omega)$$

$\rho_1(\omega) \rightarrow \infty$ for $\omega \rightarrow \Omega$ (exact excitation energy) but right-hand side remains finite for $\omega \rightarrow \Omega$

hence
$$\left(\hat{1} - \hat{\chi}_S(\omega) \left[\hat{W}_C + \hat{f}_{xc}(\omega) \right] \right) \xi(\omega) = \lambda(\omega) \xi(\omega)$$

$\lambda(\omega) \rightarrow 0$ for $\omega \rightarrow \Omega$

This condition rigorously determines the exact excitation energies, i.e.,

$$\left(\hat{1} - \hat{\chi}_S(\Omega) \left[\hat{W}_C + \hat{f}_{xc}(\Omega) \right] \right) \xi(\Omega) = 0$$

This leads to the (non-linear) eigenvalue equation

(See T. Grabo, M. Petersilka, E. K. U. G., J. Mol. Struct. (Theochem) 501, 353 (2000))

$$\sum_{q'} \left(A_{qq'}(\Omega) + \omega_q \delta_{qq'} \right) \beta_{q'} = \Omega \beta_q$$

where

$$A_{qq'} = \alpha_{q'} \int d^3 r \int d^3 r' \Phi_q(\mathbf{r}) \left(\frac{1}{|\mathbf{r} - \mathbf{r}'|} + f_{xc}(\mathbf{r}, \mathbf{r}', \Omega) \right) \Phi_{q'}(\mathbf{r}')$$

$$q = (j, a) \text{ double index}$$

$$\alpha_q = f_a - f_j$$

$$\Phi_q(\mathbf{r}) = \varphi_a^*(\mathbf{r}) \varphi_j(\mathbf{r})$$

$$\omega_q = \varepsilon_a - \varepsilon_j$$

Atom	Experimental Excitation Energies $^1S \rightarrow ^1P$ (in Ry)	KS energy differences $\Delta\epsilon_{KS}$ (Ry)	$\Delta\epsilon_{KS} + K$
Be	0.388	0.259	0.391
Mg	0.319	0.234	0.327
Ca	0.216	0.157	0.234
Zn	0.426	0.315	0.423
Sr	0.198	0.141	0.210
Cd	0.398	0.269	0.391

from: M. Petersilka, U. J. Gossmann, E.K.U.G., PRL 76, 1212 (1996)

$$\Delta E = \underbrace{\Delta\epsilon_{KS}}_{\epsilon_j - \epsilon_k} + K$$

$$K = \int d^3r \int d^3r' \varphi_j(r) \varphi_j^*(r') \varphi_k(r') \varphi_k^*(r) \left(\frac{1}{|r-r'|} + f_{xc}(r, r') \right)$$

Excitation energies of CO molecule


	State	Ω_{expt}	KS-transition	$\Delta\epsilon_{\text{KS}}$	$\Delta\epsilon_{\text{KS}} + \text{K}$
A	${}^1\Pi$	0.3127	$5\Sigma \rightarrow 2\Pi$	0.2523	0.3267
a	${}^3\Pi$	0.2323			0.2238
I	${}^1\Sigma^-$	0.3631	$1\Pi \rightarrow 2\Pi$	0.3626	0.3626
D	${}^1\Delta$	0.3759			0.3812
a'	${}^3\Sigma^+$	0.3127			0.3181
e	${}^3\Sigma^-$	0.3631			0.3626
d	${}^3\Delta$	0.3440			0.3404

T. Grabo, M. Petersilka and E.K.U. Gross, J. Mol. Struct. (Theochem) 501, 353 (2000)

approximations made: v_{xc}^{LDA} and f_{xc}^{ALDA}

Quantum defects in Helium $E_n = -\frac{1}{2(n-\mu_n)^2}$ [a.u.]

3P Series



M. Petersilka, U.J. Gossmann and E.K.U.G., in: Electronic Density Functional Theory: Recent Progress and New Directions, J.F. Dobson, G. Vignale, M.P. Das, ed(s), (Plenum, New York, 1998), p 177 - 197.




Figure 3.3: Errors of singlet excitation energies from the ground state of Be, calculated from the accurate, the OEP-SIC and x-only KLI exchange correlation potential and with different approximations for the exchange-correlation kernel (see text). The errors are given in mHartrees. To guide the eye, the errors of the discrete excitation energies were connected with lines.

(M. Petersilka, E.K.U.G., K. Burke, *Int. J. Quantum Chem.* **80**, 534 (2000))

Failures of ALDA in the linear response regime

- **H₂ dissociation is incorrect:**

$$E\left({}^1\Sigma_u^+\right) - E\left({}^1\Sigma_g^+\right) \xrightarrow{R \rightarrow \infty} 0 \text{ (in ALDA)}$$

(see: Gritsenko, van Gisbergen, Görling, Baerends, JCP 113, 8478 (2000))

- **response of long chains strongly overestimated**

(see: Champagne et al., JCP 109, 10489 (1998) and 110, 11664 (1999))


- **in periodic solids, $f_{xc}^{\text{ALDA}}(q, \omega, \rho) = c(\rho)$ whereas,**

for insulators, $f_{xc}^{\text{exact}} \xrightarrow{q \rightarrow 0} 1/q^2$ divergent.

- **charge-transfer excitations not properly described**

(see: Dreuw et al., JCP 119, 2943 (2003))

How good is ALDA for solids?



Solid Argon

L. Reining, V. Olevano, A. Rubio, G. Onida, PRL 88, 066404 (2002)

OBSERVATION:

In the long-wavelength-limit ($q = 0$), relevant for optical absorption, ALDA is not reliable. In particular, excitonic lines are completely missed. Results are very close to RPA.

EXPLANATION:

In the TDDFT response equation, the bare Coulomb interaction and the xc kernel only appear as $\text{sum}(W_C + f_{xc})$. For $q \rightarrow 0$, W_C diverges like $1/q^2$, while f_{xc} in ALDA goes to a constant. Hence results are close to $f_{xc} = 0$ (RPA) in the $q \rightarrow 0$ limit.


CONCLUSION:


Approximations for f_{xc} are needed which, for $q \rightarrow 0$, correctly diverge like $1/q^2$. Such approximations can be derived from many-body perturbation theory (see, e.g., L. Reining, V. Olevano, A. Rubio, G. Onida, PRL 88, 066404 (2002)).

WHAT ABOUT FINITE Q??


see: H.C. Weissker, J. Serrano, S. Huotari,
F. Bruneval, F. Sottile, G. Monaco, M. Krisch,
V. Olevano, L. Reining, Phys. Rev. Lett. 97,
237602 (2006)

Silicon: Loss function $\text{Im } \chi(\mathbf{q}, \omega)$





X-ray absorption spectroscopy of 3d metals



Core levels localized: TDDFT works well

**Pioneers: John Rehr
Hubert Ebert**

Detailed analysis using two-pole approximation

$$\begin{pmatrix} \omega_1 + M_{11} & M_{12} \\ M_{21} & \omega_2 + M_{22} \end{pmatrix} \beta = \Omega \beta$$

From knowledge of the KS orbitals and the KS excitation energies ω_1 , ω_2 and the experimental excitation energies Ω_1 , Ω_2 and their branching ratio, one can deduce experimental values for the matrix elements M_{ij} , i.e. one can “measure” f_{xc} .

A. Scherz, E.K.U.G., H. Appel, C. Sorg, K. Baberschke, H. Wende, K. Burke, PRL 95, 253006 (2005)